

WEST**End of Result Set**

Generate Collection

Print

L1: Entry 1 of 1

File: DWPI

Jun 15, 1999

DERWENT-ACC-NO: 1998-191484
DERWENT-WEEK: 200059
COPYRIGHT 2002 DERWENT INFORMATION LTD

TITLE: Anticancer immunoactive polysaccharide separated from
Phellinus linteus and process of making it - NoAbstract

INVENTOR: CHO, S; HAN, M ; KIM, H ; KO, K ; LEE, J ; SONG, K ; YU,
I ; YOO, I ; CHO, S M ; HAN, M W ; KIM, H M ; KOH, G S ; LEE, J H ;
SONG, G S ; YOO, I D

PATENT-ASSIGNEE:

ASSIGNEE

CODE

HANKOOK PHARM CO INC

HANKN

KOREA ADV INST SCI & TECHNOLOGY

KOAD

HAN KOOK SINYAK PHARM CO LTD

HANKN

PRIORITY-DATA: 1995KR-0029192 (September 6, 1995), 1998KR-0029082
(July 20, 1998)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
KR 197446 B1	June 15, 1999		000	C12P019/04
<u>KR 97015743 A</u>	April 28, 1997		000	C12P019/00
KR 174433 B1	February 18, 1999		000	C12P019/00

APPLICATION-DATA:

PUB-NO	APPL-DATE	APPL-NO	DESCRIPTOR
KR 197446B1	September 6, 1995	1995KR-0029192	Div ex
KR 197446B1	July 20, 1998	1998KR-0029082	
KR 97015743A	September 6, 1995	1995KR-0029192	
KR 174433B1	September 6, 1995	1995KR-0029192	

INT-CL (IPC): C12 P 19/00; C12 P 19/04

ABSTRACTED-PUB-NO: KR 97015743A

EQUIVALENT-ABSTRACTS:

CHOSEN-DRAWING: Dwg.1

TITLE-TERMS: ANTICANCER POLYSACCHARIDE SEPARATE PHELLINUS PROCESS
NOABSTRACT

DERWENT-CLASS: B04 D16

CPI-CODES: B04-C02F; B14-G03; B14-H01; D05-C08;

SECONDARY-ACC-NO:

CPI Secondary Accession Numbers: C1998-061141

대한민국 특허청 (KR)
공개특허공보 (A)

Int. Cl.
C 12 P 19 00

제 2212 호

공개일자 1997. 4. 28
출원일자 1995. 9. 6

공개번호 97-15743
출원번호 95-29192
심사청구 : 있음

발명자 유익동 대전광역시 유성구 도룡동 현대아파트 103동 401호
송봉식 대구광역시 수성구 신매동 시지 천마타운 229동 905호
이재훈 대전광역시 유성구 전민동 엑스포아파트 304동 506호
조수목 대전광역시 중구 부사동 305-2
김환모 대전광역시 유성구 덕은동 한빛아파트 133동 1301호
고병수 대전광역시 서구 삼천동 가람아파트 9동 1005호
한다우 대전광역시 서구 대동 220-2 롯데아파트 104동 1103호

출원인 한국과학기술연구원 대표자 김은영
서울특별시 성북구 하월곡동 39-1 (우: 136-130)
주최자 한국실화 대표이사 한만우
대전광역시 서구 관저동 610-7 (우: 302-243)

대리인 변리사 장상규·조현신 (전3면)

페릴리누스 린테우스 (Phellinus linteus)로부터 분리된 항암 면역활성 다당류 및 이의 제조 방법

발명자

본 발명은 페릴리누스, 린테우스 KCTC 0173BP의 균사체로부터 분리된 분자량 9,000 내지 16,000, 또는 153,000 단위의 항암 면역활성 다당류, 상기 균주의 균사체로부터 얻은 추출, 에탄올 추출 및 음이온 교환 크로마토그래피 등에 의해 분리 정제하는 것을 포함하는 상기 다당류의 제조 방법 및 상기 다당류를 활성성분으로 하는 조성물에 관한 것이다.


분리정제의 범위

1. 펠리누스 린테우스(Phellinus linteus) KCTC 0173BP 균주의 균사체로부터 분리정제된 항암 면역활성 다당류.
 2. 제1항에 있어서, 분자량이 3,000 내지 16,600달톤인 항암 면역활성 다당류.
 3. 제1항에 있어서, 분자량이 153,000달톤인 항암 면역활성 다당류.
 4. 제1항에 있어서, 상기 다당류의 당성분이 5 내지 55몰%의 글루코즈, 5 내지 30몰%의 갈락토즈, 10 내지 50몰%의 만노스, 1 내지 25몰%의 사이토스 및 5 내지 25몰%의 아라비노스로 이루어진 항암 면역활성 다당류.
 5. 펠리누스 린테우스 KCTC 0173BP 균주의 균사체를 열수 추출하고, 열수 추출물을 최종 에탄올 농도가 80%가 되도록 에탄올로 처리한 후 원심분리하여 침전물을 얻고, 침전물을 투석하고 투석물을 최종 에탄올 농도가 60%가 되도록 에탄올로 처리한 후 원심분리하여 상층액을 얻고, 상층액을 음이온 교환 크로마토그래피로 정제하는 것을 포함하는, 항암 면역활성 다당류의 제조방법.
 6. 제5항에 있어서, 상기 열수 추출시 균사체를 실온 내지 100°C의 증류수 중에서 3 내지 24시간 동안 추출하는 방법.
 7. 제1항 내지 제4항 중 어느 한 항의 항암 면역활성 다당류를 유효성분으로 포함하는 약학적 조성물.
- ※ 참고사항 : 최초출원 내용에 의하여 공개하는 것임.


도면의 간단한 설명

제1도는 펠리누스 린테우스 균주로부터 항암 면역활성 다당류를 정제하는 과정을 도시화한 것이고, 제2도는 추출물(4)의 DEAE-셀룰로즈 크로마토그래피 결과를 나타낸 것이고, 제3도는 활성물질 4-III의 겔 투과 크로마토그래피 결과를 나타낸 것이다.


제 1 도



제 2 도



제 3 도



PTO 2002-1936

Korea, Laid-open
97-15743

ANTI-CANCER IMMUNOLOGICALLY ACTIVATED POLYSACCHARIDE
SEPARATED FROM PHELLINUS LINTEUS AND ITS MANUFACTURING METHOD
[Phellinus linteus Ro Buteo Bunridoin Hangan Myeonyeok
Whalsung Tadangryu Mip Iui Jaezo Bangbeop]

Ik-Dong Yu, Kyung-Shik Song, Jae-Hoon Lee, Soc-Muk Cho,
Hwan-Muk Kim, Kyung-Soo Koh, and Man-Woo Han

UNITED STATES PATENT AND TRADEMARK OFFICE
Washington, D.C. March, 2002

Translated by: Schreiber Translations, Inc.

<u>Country</u>	:	Korea
<u>Document No.</u>	:	97-15743
<u>Document type</u>	:	Laid-open
<u>Language</u>	:	Korean
<u>Inventors</u>	:	Ik-Dong Yu, Kyung-Shik Song, Jae-Hoon Lee, Soo-Muk Cho, Hwan-Muk Kim, Kyung-Soo Koh, and Man-Woo Han
<u>Applicants</u>	:	Korea Institute of Science and Technology Korea Shinyak Co., Ltd.
<u>IPC</u>	:	C 12 P 19/00
<u>Application date</u>	:	September 6, 1995
<u>Publication date</u>	:	April 28, 1997
<u>Foreign Language Title</u>	:	Phellinus linteus Ro Buteo Bunridoin Hangam Myeonyeok Whalsung Tadangryu Mip Iui Jaezo Bangbeop
<u>English Title</u>	:	ANTI-CANCER IMMUNOLOGICALLY ACTIVATED POLYSACCHARIDE SEPARATED FROM PHELLINUS LINTEUS AND ITS MANUFACTURING METHOD

Abstract

/1*

The present invention pertains to an anti-cancer immunologically activated polysaccharide with a molecular weight of 9,000-16,000 or 153,000 dalton separated from a mycelium of *Phellinus linteus* KCTC 0173BP, a method for manufacturing the above-mentioned polysaccharide including a hydrothermal extraction from the mycelium of the above-mentioned strain, ethanol extraction, and separation and purification by anion exchange chromatography, etc., and a pharmaceutical composition containing the above-mentioned polysaccharide as an active ingredient.

*Numbers in the margin indicate pagination in the foreign text.

1. Title of the Invention: ANTI-CANCER IMMUNOLOGICALLY ACTIVATED
POLYSACCHARIDE SEPARATED FROM
PHELLINUS LINTEUS AND ITS
MANUFACTURING METHOD

2. Claims

1. An anti-cancer immunologically activated polysaccharide characterized by being separated and purified from a mycelium of Phellinus linteus KCTC 0173BP strain.

2. The anti-cancer immunologically activated polysaccharide of Claim 1 characterized by the fact that the molecular weight is 9,000-16,000 dalton.

3. The anti-cancer immunologically activated polysaccharide of Claim 1 characterized by the fact that the molecular weight is 153,000 dalton.

4. The anti-cancer immunologically activated polysaccharide of Claim 1 characterized by the fact that the sugar component of the above-mentioned polysaccharide is composed of 5-55 mol% glucose, 5-30 mol% galactose, 10-50 mol% mannose, 1-25 mol% xylose, and 5-25 mol% arabinose.

5. A method for manufacturing an anti-cancer immunologically activated polysaccharide characterized by that a mycelium of Phellinus linteus KCTC 0173BP strain is hydrothermally extracted; the

hydrothermal extract is treated with ethanol so that the final ethanol concentration may be 80% and is then centrifuged, so that a precipitate is obtained; the precipitate is dialyzed; the dialysate is treated with ethanol so that the final ethanol concentration may be 60% and is then centrifuged, so that a supernatant is obtained; and the supernatant is purified by an anion exchange chromatography.

6. The method of Claim 5 characterized by the fact that in the above-mentioned hydrothermal extraction, the mycelium is extracted for 3-24 h in distilled water at room temperature to 100°C.

7. A pharmaceutical composition, characterized by including the anti-cancer immunologically activated polysaccharide of any of Claims 1-4 as an active ingredient.

* Remarks: Initially disclosed according to the initial filing.

3. Brief description of the figures

Figure 1 is a diagram showing a process for purifying the anti-cancer immunologically activated polysaccharide from *Phellinus linteus* strain. Figure 2 shows the results of a DEAE-cellulose chromatography of a coarse extract (4). Figure 3 shows the results of a gel permeation chromatography of an activator 4-III.

8. Coarse extract (3) of dialysis membrane inside
9. Extract of dialysis membrane outside
10. Ethanol treatment (final concentration 60%)
Centrifuging at 12,000 rpm for 30 min
11. Supernatant
12. Freeze-drying
13. Precipitate
14. Freeze-drying
15. 80% ethanol undissolved, 60% ethanol dissolved
Coarse extract (4)
16. 80% ethanol and 60% ethanol undissolved
Coarse extract (5)
17. DEAE cellulose
Ion chromatography
18. Elution with a buffer solution
Freeze-drying
19. 0.1-1 M NaCl concentration gradient elution
Dialysis, freeze-drying
20. HW 65F
Gel permeation chromatography




Figure 2:

1. Carbohydrate/A490
2. Fraction No.
3. Protein/A750
4. Carbohydrate/A490
5. Protein/A750




Figure 3:

1. Sugar A490
2. Fraction No. (4 ml/tube)
3. Protein A595